Toward the Development of Severe Weather Outlooks

Scott Weaver NOAA Climate Prediction Center



Community White Paper & NOAA Fact Sheet

Advancing the Nation's capability to anticipate tornado and severe weather risk

Scott Weaver (NOAA/CPC), Jeff Trapp (Purdue University), Michael K. Tippett (IRI), Russell Schneider (NOAA/SPC), Phil Pegion (NOAA/ESRL,CIRES), Sang-Ki Lee (NOAA/AOML), Wavne Higgins (NOAA/CPC), Andy Dean (NOAA/SPC), Greg Carbin (NOAA/SPC), Harold Brooks (NOAA/NSSL), Mike Baldwin (Purdue University), Francisco Alvarez (Saint Louis

forecast models; their ongoing work similarly demonstrates the viability of using global climate models, including the CFS, in such a dynamical downscaling approach. The forecast model - Weather Research and Forecasting (WRF) model - is configured such that convective clouds and storms are explicitly represented over the entire The output of such domain. convection-permitting WRF-model integrations over months and seasons is then mined to determine the spatial distribution and frequency of convective storms by type and severity. Figure 3 shows an example



Fig 3. Example of predicted storm occurrence (simulated radar reflectivity : predicted severe storm 40 dBZ) over the period April, May, Jun, 2012, generated by WRF enabled occurrence over the period April downscaling of the CFS model.

through June of 2012, generated by WRF-enabled downscaling of the CFS model.

Observational Databases

Underpinning our current understanding of the severe weather environment and its climate connections is a long term (1950-present) historical database of F-scale tornado counts. Since this database was not intended to be a consistent homogenous long-term climate record of tornadic and severe weather parameters, there are inherent inconsistencies as a result of public awareness, tornado reporting practices, NWS guidelines, and other sources of inhomogeneity. These issues may introduce spurious

trends in the long-term tornado data. However, it has been demonstrated that much of this trend can be ameliorated by focusing on the F1-F5 tornado counts only (Verbout et al. 2006) as demonstrated in Figure 4 which shows that much of the trend can be explained through the timeseries evolution of the F0 tornado counts. Nevertheless, it is necessary to explore other novel ways to further homogenize the long-term historical tornado database while simultaneously taking



State of the Science FACT SHEET

Tornadoes, Climate Variability, and Climate Change

This assessment of tornado activity and climate was developed by scientists and communication experts from the National Oceanic and Atmospheric Administration (NOAA)

Tornadoes are intense rotating vertical columns of air that pose a great threat to lives and property. They typically form in an environment where winds are rapidly changing direction and speed with height (commonly referred to as wind shear) and the atmosphere is convectively unstable Tornado strength is classified according to the Fujita (F) Scale F0-F5, with F0 being the weakest and F5 the strongest. Tornadic activity refers to the number and intensity of tornadoes over a given region, season, or year. While tornadoes can occur during any season in the U.S., they are most likely during the spring months of March, April, and May.

Given the right set of atmospheric conditions, tornadoes can occur almost anywhere. However, the areas of the U.S. most susceptible include the Great Plains, Midwest and South. The configuration of the topography of the North American continent (Rocky Mountains, Great Plains, and proximity to the Gulf of Mexico) contributes to the development of large-scale weather systems capable of supporting severe thunderstorms and related tornado

Are the frequency and/or intensity of tornadoes increasing?

Underpinning our current understanding of tornado activity is a long-term (1954-present) record of historical tornado counts from NOAA's Severe Weather Database (SWD). Given that the SWD was not intended to be a consistent homogenous long-term climate record of tornadoes, there are inconsistencies over time as a result of changes in public awareness, tornado reporting practices, Doppler radar technology, and National Weather Service (NWS) guidelines, to name a few.

These inconsistencies have likely introduced artificial trends in the long-term tornado data making attribution of long-term changes in tornado frequency/intensity difficult to determine. This issue is highlighted by a comparison of all tornado counts (F0-F5) with only the F1-F5 tornadoes (Figure 1). Removing the F0 counts from the database nearly eliminates the trend. Despite the potential for spurious trends in the SWD, this does not rule out the possibility that a portion of the trend is due to climate change or natural climate variability

*A new Enhanced Fulita scale was adopted in 2007.

What is the role of natural climate variability in tornadic activity?

Emerging evidence suggests that natural climate variations such as El Nino and La Nina events and in narticular the transition from one to another in spring have the potential to modulate the environmental factors

SWD: Number of U.S. Tornadows (Annual)





Figure 1. The number of F0-F5 (upper), F1-F5 (lower) annual tornado counts for 1954-2011 from the NOAA Severe Weather Database. The grav lines show the linear trend for the respective F-scale intervals. (From Lee et al. 2012)

conducive to tornado formation¹. Natural variations acting on decadal timescales (e.g., Atlantic Multidecadal and Pacific Decadal Oscillations) have also been similarly implicated. However, uncertainty is higher because these phenomena are influential over the course of roughly 30 vears and the tornado record is only approximately 60 years long. The link between these inter-annual-todecadal natural climate variability modes and variations in tornado activity is typically via shifts in the large scale upper and lower level jet streams across North America that act as focusing mechanisms for severe thunderstorm development. Shifts in the placement and strength of these jet streams will influence the locations of

April 2013

NOAA.gov

Can we predict long-term changes in tornado activity?

Emerging evidence suggests it may be possible to provide prediction of overall tornadic activity over the U.S.³ However, the viability of providing long-term (i.e., monthly and seasonal) outlooks of tornadoes has not yet reached a level necessary to provide actionable

atmospheric instability and wind shear that promote enhanced or reduced tornado activity

Does climate change impact tornado activity?

Despite the existence of artificial trends in the SWD there is still a potential role for changes in tornado activity due to climate change. Given that tornado activity is dependent upon two things: the strength of atmospheric instability, which promotes rising air and thunderstorm formation; and vertical wind shear, which provides the necessary rotation for tornadic thunderstorms (Figure 2). the best approach is to assess the likely impact climate change will have on these proxy indicators. Nevertheless. this is not straightforward since early indications show a likely tug-of-war between these two mechanism

In general, climate change is expected to increase atmospheric instability by increasing temperature and humidity in the lower atmosphere, while simultaneously weakening vertical wind shear through a reduction in the surface pole-to-Equator temperature gradient. The result is regions that currently see the highest frequency of tornado activity may experience reductions due to weakened vertical wind shear, unless this is offset by the increases in atmospheric instability. Additionally, these factors could modify the seasons of tornado activity, i.e., enhance cool season activity. Some observations indicate storminess is increasing in general, suggesting atmospheric instability may be winning out over the wind shear influence.



how instability and wind shear influence unstable rising air (red arrows) is forced to rotate by the increased wind speed between the lower (green) and mid-toupper levels (purple-orange).

Does global warming influence tomado activity? EOS, Trans. Amer. Geophys. Union, 89, 53, doi:10.1029/2008EO530001. ³ Tippett, M. K., A. H. Sobel, and S. J. Camargo, 2012:

Association of U.S. tornado occurrence with monthly environmental parameters. Geophys. Res. Lett., 39, L02801. doi:10.1029/2011GL050368.

information. Since current versions of climate models do

not have the resolution necessary for explicitly

representing the extremely small-scale tornadic

structures, other techniques are being explored. The

current strategy focuses on the prediction of local proxy

indicators (atmospheric instability and wind shear) and remote climate signals (El Nino/La Nina transitions and

other climate variability modes). An effort is underway to

develop a quantitative monthly-to-seasonal prediction system that exploits new insights and knowledge as they

The potential for predicting longer-term tornado activity

(i.e., decadal-to-centennial) is much less clear given the

inherent uncertainties in the model based prediction of

decadat-to-centennial regional climate change. To be sure, there is much room for improvement and large gaps

it is widely believed among the scientific community there

How should research be directed to improve

Near-term research should target several crucial areas:

• Improve understanding of the influence of large-scale

· Improve understanding of potential climate change

impacts on the frequency and/or intensity of tornadoes

Explore new ways to make better use of the historical

database (i.e., focusing on tornado days, path length

and width, etc.) and develop a modern record that

Integrate societal needs into the research process to

Lee, S-K. R. Atlas, D. Enfield, C. Wang, and H. Liu,

2012: Is there an optimal pattern that enhances large-

² Diffenbaugh, N. S., R. J. Trapp, and H. Brooks, 2008:

scale atmospheric processes conducive to outbreaks in the U.S.? J. Climate, 26, 1626-1642.

better align the applicability of research advances.

natural climate variations on tornadic activity.

Advance modeling and downscaling strategies

exploits recent technological advances

Additional Resources

References:

exist in current understanding on all timescales. Ho

is great potential for research to fill these gaps.

are discovered

understanding?

April 2013

es conducive to tornado



Tornadoes 2005-14



How can we improve understanding of long term changes and year-toyear variability?









Combined Influence of NALLJ Modes 1 & 2 On Regional Tornadic Activity



Pattern Correlations Between CFSv2 (172 runs) Loading Patterns and CFSR

EOF-1

EOF-2







CFSv2 0-month lead 20-member forecast of AMJ NALLJ Mode 2

Regional Tornadic Variability SST

- SST linkages may offer prediction opportunity.
- Compare and contrast the spatial patterns of global SST variability to regional tornado indices.
- Strategy exhibits no a priori assumption regarding the structure of associated SST variability, the case and limitation when targeting connectivity to indices of ENSO.
- Northern Great Plains & Southeast example.



160W

120W

80W

40W

160E

405 + 40E

80E

120E



AMJ NP DAYS Tornado SST Correlation





A Prediction Example



U.S. Climate Regions

Tornado Index Regional Correlations

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep		Nov	Dec
South	0.16	0.36	0.29	0.05	0.28	0.51	0.09	0.34	0.05	0.30	0.29	0.33
Southeast	0.22	0.24	0.00	0.41	0.66	0.25	-0.01	0.00	0.49	0.26	0.45	0.47
Central	0.47	0.50	0.64	0.23	0.37	0.45	0.42	0.05	0.19	0.03	0.24	0.42
Midwest			-0.12	0.58	0.15	0.67	0.39	0.42	0.02	0.39	-0.04	
Plains			0.12	0.37	0.40	0.50	0.53	0.27	-0.03	0.03		
Northeast				0.15	0.05	0.15	0.41	0.18	0.70	0.15	-0.02	
Southwest				0.02	-0.10	0.32	0.04	-0.01	-0.44	0.30		
Northwest				-0.14	0.15	0.30		0.19				
West		0.21	0.34	0.13								

TABLE 4. Correlation between the index and reported number of tornadoes by U.S. climate region and month for the period 1979-2010. Significant correlations are in bold font. Regions and months with less than 32 reported tornadoes are omitted.

Mike Tipett, IRI, 2013

A Final Word

These are just a few vignettes. Other topics include:

- Downscaling methodologies.
- Intraseasonal variability. Some links to MJO in spring.
- New ways to correct the historical database.
- Exploiting modern technology to develop and maintain a new climate record for the next generation of scientists.
- White Paper available upon request or download at:

www.cpc.ncep.noaa.gov/products/severe_weather/ Climate.Severe.Weather.White.Paper.Apr2013.pdf







