

Numerical modeling studies on the influence of the Arctic on the atmospheric large-scale circulation of the Northern mid-latitudes

Question

- What happens to the weather and climate of the Northern mid-latitudes if the sea ice and the Arctic atmosphere change faster than anticipated?
- Idealized model studies which *only* consider the influence of the Arctic and keep the influence of the mid-latitudes and tropics as small as possible

Experiments

- Atmosphere-only relaxation experiments (14 days)
- Idealized atmosphere-only experiments with reduced sea ice thickness (15 days, some 90 days)
- Idealized coupled experiments with initially reduced sea ice thickness (1 year)
- Idealized coupled experiments with modified albedo, lead closing parameter, longwave radiation (150 years)

Atmosphere-only relaxation experiments




Fig. 1: Forecast error reduction (%) through relaxation of prognostic variables north of 75°N in winter
Within Northern mid-latitudes Northern Asia most affected – due to northerly component in mean westerly flow




Fig. 2: 2 m temperature anomaly (K) in cases of strongly improved forecasts in Northern Asia winter
Link to European cold winters
Semmler et al., 2017, submitted to AAS
Jung et al., 2014, GRL

Idealized atmosphere-only experiments




Fig. 3: Mean vertical temperature profiles for CTL (black contour lines), interval 4 (°C), and differences (colour shading (K)) between ice-reduced (RED) and CTL
Semmler et al., 2016a
Mainly boundary layer affected

Idealized short coupled experiments




Fig. 4: Synoptic activity and Eady growth rate in CTL simulation and difference ice-reduced (RED) minus CTL
Semmler et al., 2016b
Less synoptic activity but stronger Eady growth rate in Arctic, southward shift of storm track

Idealized long coupled experiments




Fig. 5: Mean sea surface height response in the last 60 years of the 150 year-simulations
Spin-up of the Beaufort Gyre: less and/or thinner sea ice cover permit stronger momentum flux into the ocean
Pathway shift of mid-latitude surface currents: negative AO forces southward shift of Western Boundary Current Extensions, consequence: warmer and saltier North Atlantic current
Campos et al., 2017, in prep.

Conclusions

- Reduced sea ice increases temperature mainly in Arctic boundary layer
- Strongest pathway from Arctic to Northern mid-latitudes: Barents Sea / Kara Sea area -> Siberia
- Reduced westerly flow especially over Eurasian sector along with some cooling
- Less synoptic activity but stronger Eady growth rate in the Arctic (vertical stability increase not as relevant as vertical wind shear decrease)
- Southward atmospheric storm track shift
- Encouraging: results consistent between different methods and different time scales
- In long coupled simulations southward atmospheric storm track shift reflected in the ocean. Generally more active ocean circulation in Arctic and sub-Arctic.

References:

- Jung, T., Kasper, M.A., Semmler, T., Serrar, S. (2014): Arctic influence on subseasonal midlatitude prediction. *Geophysical Research Letters*, 41, 3676-3680, doi: 10.1002/2014GL059961
- Semmler, T., Jung, T., Serrar, S. (2016a): Fast atmospheric response to a sudden thinning of Arctic sea ice. *Clim. Dyn.*, 46, 1015, doi: 10.1007/s00382-015-2629-7

- Semmler, T., Stulic, L., Jung, T., Tilinina, N., Campos, C., Gulev, S., Koracin, D. (2016b): Seasonal Atmospheric Responses to Reduced Arctic Sea Ice in an Ensemble of Coupled Model Simulations. *Journal of Climate*, 29, 5893-5913, doi: 10.1175/JCLI-D-15-0586.1
- Semmler, T., Jung, T., Kasper, M.A., Serrar, S. (2017): Using NWP to assess the influence of the Arctic atmosphere on mid-latitude weather and climate. *Submitted to Advances in Atmospheric Sciences*.
- Campos, C., Semmler, T., Jung, T. (2017): The response of Northern hemisphere oceans to the Arctic sea ice decline. *In preparation*.