

The Seasonal Footprinting Mechanism in CFSv2: Simulation and Impact on ENSO Prediction

^{1,2}Kathy Pegion and ¹Micheal Alexander

¹NOAA/Earth System Research Laboratory/Physical Sciences Division

²CIRES/University of Colorado

The seasonal footprinting mechanism (SFM) is thought to be a pre-cursor to the El Nino Southern Oscillation (ENSO). Fluctuations in the North Pacific Oscillation (NPO) impact the ocean via surface heat fluxes during winter, leaving a sea-surface temperature (SST) “footprint” in the subtropics. This footprint persists through the spring, impacting the tropical Pacific atmosphere-ocean circulation throughout the following year.

The simulation of the SFM in the NCEP/Climate Forecast System, version 2 (CFSv2) is likely to have an impact on operational predictions of ENSO. We explore the simulation of the SFM in the CFSv2 decadal simulations and compare with reanalysis products. Additionally, we investigate the relationship between the SFM and ENSO prediction skill in the NCEP/CFSv2 retrospective forecasts.

Results indicate that the CFSv2 has a weak simulation of the NPO with the southern lobe of the NPO located too far to the north. As a result, the SFM in the CFSv2 is not as effective at impacting the Tropics as indicated by reanalysis. This leads to less reliable predictions of ENSO following a positive or negative NPO event than when the NPO is not active.